Synthetic Communications

An International Journal for Rapid Communication of Synthetic Organic Chemistry

ISSN: 0039-7911 (Print) 1532-2432 (Online) Journal homepage: <http://www.tandfonline.com/loi/lsyc20>

Multicomponent reaction of benzyl halides: Synthesis of [1,2,4]triazolo/benzimidazolo quinazolinones

Mallappa Beerappa & Kalegowda Shivashankar

To cite this article: Mallappa Beerappa & Kalegowda Shivashankar (2016) Multicomponent reaction of benzyl halides: Synthesis of [1,2,4]triazolo/benzimidazolo quinazolinones, Synthetic Communications, 46:5, 421-432

To link to this article: <http://dx.doi.org/10.1080/00397911.2016.1140784>

View supplementary material [↗](#)

Accepted author version posted online: 20 Jan 2016.
Published online: 20 Jan 2016.

Submit your article to this journal [↗](#)

Article views: 156

View related articles [↗](#)

View Crossmark data [↗](#)

Citing articles: 2 View citing articles [↗](#)

Multicomponent reaction of benzyl halides: Synthesis of [1,2,4]triazolo/benzimidazolo quinazolinones

Mallappa Beerappa and Kalegowda Shivashankar

P.G. Department of Chemistry, Central College Campus, Bangalore University, Bangalore, Karnataka, India

ABSTRACT

We demonstrated a simple, highly efficient, one-pot method to construct triazolo/benzimidazolo quinazolinones through a cascade of oxidation, Knoevenagel condensation, and Michael addition followed by cyclization and dehydration. This protocol tolerates easily available benzyl halides, 2-amino benzimidazole/3-amino-1,2,4-triazole, and α -hydroxy C-H acids as starting materials using trimethyl amine *N*-oxide in ethanol. To the best of our knowledge, this is the first example of synthesis of triazolo/benzimidazolo quinazolinones directly from benzyl halides in one pot. Simple procedure, environmental friendliness, mild reaction conditions, and good yields are the attractive features of this method.

GRAPHICAL ABSTRACT

ARTICLE HISTORY

Received 16 October 2015

KEYWORDS

Benzimidazole;
multicomponent reaction;
quinazolinone; triazole;
trimethyl amine *N*-oxide

Introduction

Multicomponent reactions (MCRs) have been refined in recent years as powerful and necessary tools in synthetic organic and medicinal chemistry. This powerful tool combines both the environmental and economic aspects of organic chemistry that are essential for industrial and academic applications. MCRs offer remarkable advantages such as convergence, reduction in the amount of workup, operational simplicity in the separation and purification processes, minimal waste production, and green transformations.^[1–5]

The synthesis of quinazolinone analogs is a significant synthetic reaction in organic chemistry as these scaffolds are found to be very important core structures in various natural products^[6,7] and a wide range of biologically active synthetic compounds.^[8,9] Many

CONTACT K. Shivashankar ✉ shivashankark@gmail.com P.G. Department of Chemistry, Central College Campus, Bangalore University, Bangalore, Karnataka, India.

Color versions of one or more of the figures in this article can be found online at www.tandfonline.com/lsyc.

Supplemental data for this article can be accessed on the [publisher's website](#).

© 2016 Taylor & Francis

Figure 1. Quinazolinone-derived commercially available drugs.

commercially available drugs (Fig. 1) including ispinesib (anticancer), dacomitinib (anticancer), methaqualone (antimalarial), albaconazole (antifungal activity), balaglitazone (antidiabetic and hypolipidemic property), and anagrelide (for the treatment of essential thrombocythosis) are derived from quinazolinone core entities.^[10] Quinazolinone moieties exhibit a variety of biological activities such as antitumor,^[11] antiviral,^[12] anticancer,^[13] antileishmanial,^[14] antimicrobial,^[15] anti-inflammatory,^[16] antitubercular,^[17] antimalarial,^[18] antifungal,^[19] and antibiotic.^[20] Hence, the synthesis of quinazolinone has evoked much interest, as a result of which a variety of synthetic methodologies have been reported.^[21–23] The most important approaches are (i) one-pot condensation of aromatic aldehydes, 2-amino benzimidazole/3-amino-1,2,4-triazole, and α -hydroxy C-H acids,^[24] (ii) three-component reaction of 2-amino benzamides, aryl halides, and isocyanides,^[25] (iii) one-pot reaction with *o*-amino benzamides and primary alcohols,^[26] (iv) cascade synthesis of phenylamino substituted quinazolinones from urea derivatives and *tert*-butyl isocyanide,^[27] (v) oxidative radical skeletal rearrangement induced by oxygen,^[28] (vi) synthesis of quinazolinone and quinazoline from aerobic oxidative C(sp³)–H amination/C–N cleavage.^[29] Other methods have also been developed within the past three decades.^[30] In the past years, a few methods have described the one-pot multicomponent synthesis of quinazolinones based on catalysts such as $\text{NH}_2\text{SO}_3\text{H}$,^[31] thiamine hydrochloride,^[32] CuCl ,^[33] chitosan,^[34] nafen-H,^[35] Mg-Al-CO_3 ,^[36] silica gel,^[37] and $\text{H}_6\text{P}_2\text{W}_{18}\text{O}_{60} \cdot 18\text{H}_2\text{O}$ ^[38] from the reaction of aromatic aldehydes, 2-amino benzimidazole/3-amino-1,2,4-triazole, and α -hydroxy C-H acids. Nevertheless, these methods may be restricted to benzaldehyde and have some significant limitations, such as harsh reaction conditions, long reaction time, tedious workup, poor yields, and use of metal- and moisture-sensitive catalysts. Hence, the development of a simple, high-yielding, and environmentally kind procedure for the one-pot multicomponent synthesis of quinazolinone analogs is still reasonable.

As part of our continuing efforts on the development of new methods for the construction of biologically active compounds,^[39] we have reported synthesis of

pyran-based heterocycles via one-pot reaction of benzyl halides, α -hydroxy C-H acids, and active methylene compounds.^[40] Herein, we investigated a new and straightforward approach for the synthesis of triazolo/benzimidazolo quinazolinones that involved an in situ oxidation of benzyl halides with trimethyl amine *N*-oxide into corresponding aldehydes, which they undergo a three-component reaction with α -hydroxy C-H acids and 2-aminobenzimidazole/3-amino-1,2,4-triazole without addition of any other catalyst. The wider scope and synthetic utility of trimethyl amine *N*-oxide for the synthesis of heterocyclic compounds in one pot has not been explored even though this reagent has been used as an oxidizing agent.^[41,42] This procedure developed the known reaction using the novel methodology, while not limiting one of the substrates to an aldehyde.^[43]

Results and discussion

To set up the optimal reaction conditions for the construction of quinazolinone analogs, our preliminary experiments were focused on a one-pot reaction involving 3-cyanobenzyl bromide (1 equiv), 2-amino benzimidazole (1 equiv), and dimidone (1 equiv) in the presence of trimethyl amine *N*-oxide (2 equiv) as an oxidizing agent. For a low-hazard and ecofriendly process, ethanol is used as a preferred solvent for all the optimization experiments. The reactions were performed with heating at 70 °C. The yield of the product was relatively superior. However, the yield decreased when the reaction proceeded to room temperature or more than 70 °C. Based on these results, a reaction of 3-cyanobenzyl bromide (1 equiv), 2-amino benzimidazole (1 equiv), and dimidone (1 equiv) in the presence of trimethyl amine *N*-oxide (2 equiv) stirring in ethanol at 70 °C was considered as a standard and model reaction (Scheme 1) to optimize the reaction conditions for the one-pot synthesis of other triazolo/benzimidazolo quinazolinones.^[44]

To evaluate the effect of organic solvents to optimize the reaction conditions using trimethyl amine *N*-oxide as an oxidizing agent for the synthesis of quinazolinone analogs (Table 1), ethanol was replaced with other polar solvents such as tetrahydrofuran (THF), acetonitrile, dimethylformamide (DMF), water, acetic acid, and methanol (entries 4 and 6–10). The results showed that using ethanol as the solvent gave the best yield of the respective product with greater reaction rate (entry 5). The use of nonpolar solvents such as toluene, chloroform, and dioxane afforded the desired product in traces (entries 1–3). Therefore, ethanol was selected as a suitable medium to run the further reaction.

The scope of the present protocol was further investigated by the replacement of dimidone with ethylacetoacetate. To our observation, the reaction underwent successful

Scheme 1. Synthesis of 12(3-cyanophenyl)-3,3-dimethyl-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-b]quinazolin-1-one (4a).

Table 1. Optimization of solvents for the synthesis (**4a**).

Entry	Solvent	Time (min)	Yield (%)
1	Toluene	95	Trace
2	CHCl ₃	95	Trace
3	Dioxane	90	Trace
4	THF	80	55
5	Ethanol	70	96
6	Acetonitrile	75	86
7	DMF	80	78
8	Water	80	68
9	Acetic acid	80	65
10	Methanol	70	70

condensation under similar reaction conditions to give the corresponding quinazolinone analogs in excellent yields (Table 2, entries 8 and 9). Reactions were also carried out by replacing 2-amino benzimidazole with 3-amino-1,2,4-triazole. The reaction underwent smooth cyclization to give respective quinazolinone derivatives in virtually good yields (Table 2, entries 10–14). Thereafter, a series of different substituted quinazolinone analogs were synthesized from different benzyl halides containing electron-withdrawing and electron-donating groups using trimethyl amine *N*-oxide in ethanol at 70 °C in good yields^[45] (Table 2). In general, the overall yield ranged from 96% for 3,3-dimethyl-12(4-nitro-phenyl)-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-*b*]quinazolin-1-one (**4h**) (entry 8) to 85% for 6,6-dimethyl-9-(4-hydroxy-phenyl)-5,6,7,9-tetrahydro-4*H*-1,2,4-triazolo[5,1-*b*]quinazolin-8-one (**4m**) (entry 13).

The proposed mechanism (Scheme 2) is similar to the well-known mechanism reported in the literature.^[46–49] In the first step, oxidation proceeds via nucleophilic substitution of the benzyl halides with the oxygen of the trimethyl amine *N*-oxide to give a *N*-alkoxy salt. Afterward, proton elimination at the carbon carrying the oxygen leads to the formation of an aldehyde, giving trimethyl amine hydrogen halide salt, which in the second step would be able to acid catalyze Knoevenagel condensation between aldehyde and α -hydroxy C-H acid to give the heterodiene. The third step is the Michael addition of the amino compound on the heterodiene, followed by cyclization and dehydration in the presence of acid to afford the desired product.

Conclusion

In conclusion, we have developed a new and straightforward methodology for the one-pot synthesis of quinazolinone analogs via trimethyl amine *N*-oxide-catalyzed in situ oxidation of benzyl halides into benzaldehydes followed by the three-component reaction with α -hydroxy C-H acids and amine compounds. Mild reaction condition, one-pot operation, easy isolation of product, and broad functional group tolerance are the main advantages of this reaction.

Experimental

All the reagents were purchased commercially and were used directly without further purification. The melting points were measured by the open capillary tubes using electric melting-point equipment and are uncorrected. The infrared (IR) spectra were obtained using an Agilent Cary 630 FT-IR spectrophotometer. ¹H NMR (400 MHz)

Table 2. Synthesis of quinazolinone analogs (4a–n).

Entry	Benzyl halide	α -Hydroxy C-H acid	Amine source	Quinazolinone	Time (min)	Yield (%)	Obs. mp (°C)	Lit. mp (°C) ^{Ref}
1					70	94	>300	
2					70	92	>300	
3					70	92	>300	>300 ^[31]

(Continued)

Table 2. Continued.

Entry	Benzyl halide	α -Hydroxy C-H acid	Amine source	Quinazolinone	Time (min)	Yield (%)	Obs. mp ($^{\circ}$ C)	Lit. mp ($^{\circ}$ C) ^{Ref}
4					75	90	>300	>300 ^[31]
5					70	92	>300	>300 ^[31]
6					85	88	>300	>300 ^[50]

(Continued)

Table 2. Continued.

Entry	Benzyl halide	α -Hydroxy C-H acid	Amine source	Quinazolinone	Time (min)	Yield (%)	Obs. mp ($^{\circ}$ C)	Lit. mp ($^{\circ}$ C) ^{Ref}
7					85	88	>300	
8					70	96	>300	>300 ^[51]
9					70	92	>300	

(Continued)

Table 2. Continued.

Entry	Benzyl halide	α -Hydroxy C-H acid	Amine source	Quinazolinone	Time (min)	Yield (%)	Obs. mp ($^{\circ}$ C)	Lit. mp ($^{\circ}$ C) ^{Ref}
10					65	94	>300	
11					70	92	292	
12					70	92	310	303–305 ^[31]

(Continued)

Table 2. Continued.

Entry	Benzyl halide	α -Hydroxy C-H acid	Amine source	Quinazolinone	Time (min)	Yield (%)	Obs. mp ($^{\circ}$ C)	Lit. mp ($^{\circ}$ C) ^{Ref}
13					80	85	>300	>300 ^[31]
14					65	94	>300	307–309 ^[31]

Scheme 2. Plausible reaction mechanism for the formation of quinazolinone derivatives.

and ^{13}C NMR (100 MHz) spectra were recorded on a Bruker spectrometer using dimethylsulfoxide ($\text{DMSO}-d_6$) as solvents and tetramethylsilane (TMS) as an internal standard at IISC, Bangalore. The chemical shifts were given on the δ scale (ppm). The purity of the compounds was checked by thin-layer chromatography (TLC, Merck silica-gel 60 F254 precoated aluminium sheets). The CHN analysis was performed on an Elemental Vario Micro Cube CHN rapid analyzer. All the synthesized compounds gave agreeable elemental study results.

Typical procedure for the synthesis of 12(3-cyanophenyl)-3,3-dimethyl-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-b]quinazolin-1-one (4a)

Trimethyl amine *N*-oxide (290 mg, 3.0 mmol) was added to a solution of 3-cyanobenzyl bromide (290 mg, 1.5 mmol), dimedone (210 mg, 1.5 mmol), and 2-aminobenzimidazole (200 mg, 1.5 mmol) in ethanol (10 mL) in a round-bottomed flask. The resulting reaction mixture was heated at 70°C in an oil bath for 70 min. The progress of the reaction was monitored by TLC. After completion of the reaction, the reaction mixture was allowed to cool to room temperature. The solid was collected by filtration at pump, washed with ethanol, and recrystallized from hot ethanol to obtain pure product.

12(3-Cyano-phenyl)-3,3-dimethyl-1,2,3,4,5,12-hexahydrobenzo[4,5]imidazo[2,1-b]quinazolin-1-one (4a)

Yield 94%; white solid; mp $> 300^\circ\text{C}$; IR (ATR, cm^{-1}): 3394 (NH), 2018 (CN), 1660 ($\text{C}=\text{O}$) cm^{-1} ; ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 0.98 (s, 3H, $-\text{CH}_3$), 1.04 (s, 3H, $-\text{CH}_3$),

2.22 (m, $J=4.0$ Hz, 2H, $-\text{CH}_2$), 2.48–2.64 (m, $J=4.0$ Hz, 2H, $-\text{CH}_2$), 6.50 (s, 1H, CH), 6.94–7.92 (m, 8H, Ar-H), 11.18 (s, 1H, NH) ppm; ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 26.7, 28.1, 28.4, 32.2, 49.7, 53.6, 105.3, 109.9, 111.0, 117.0, 118.5, 120.7, 122.0, 129.8, 130.9, 131.5, 141.8, 142.8, 145.0, 150.9, 158.1, 159.1, 192.6 ppm. Anal. calcd. for $\text{C}_{23}\text{H}_{20}\text{N}_4\text{O}$: C, 74.98; H, 5.47; N, 15.21. Found: C, 74.82; H, 5.31; N, 15.13.

Funding

We acknowledge a grant from the Department of Science and Technology–Science Engineering Research Board (DST-SERB), New Delhi, India, under Fast Track Young Scientist Scheme [Grant No. SB/FT/CS-028/2013].

References

- [1] Gu, Y. *Green Chem.* **2012**, *14*, 2091.
- [2] Wei, E.; Liu, B.; Lin, S.; Liang, F. *Org. Biomol. Chem.* **2014**, *12*, 6389.
- [3] Nourisefat, M.; Panahi, F.; Nezhad, A. K. *Org. Biomol. Chem.* **2014**, *12*, 9419.
- [4] Shivashankar, K.; Shastri, L. A.; Kulkarni, M. V.; Rasal, V. P.; Saindane, D. M. *J. Indian Chem. Soc.* **2009**, *86*, 265.
- [5] Pereira, M. F.; Thiery, V. *Org. Lett.* **2012**, *14*, 4754.
- [6] Liu, F.; Kaselj, M.; Isome, Y.; Ye, P.; Sargent, K.; Sprague, K.; Cherrak, D.; Wilson, C. J.; Si, Y.; Yohannes, D.; Ng, S. C. *J. Comb. Chem.* **2006**, *8*, 7.
- [7] Sawant, S. D.; Srinivas, M.; Reddy, G. L.; Rao, V. V.; Singh, P. P.; Vishwakarma, R. A. *Tetrahedron Lett.* **2012**, *53*, 6195.
- [8] Khan, I.; Ibrar, A.; Abbas, N.; Saeed, A. *Eur. J. Med. Chem.* **2014**, *76*, 193.
- [9] Rajput, R.; Mishra, A. P. *Int. J. Pharm. Sci.* **2012**, *4*, 66.
- [10] (a) Vijayakumar, B.; Prasanthi, P.; Teja, K. M.; Reddy, K. M. K.; Nishanthi, P.; Nagendramma, M.; Nishanthi, M. *Int. J. Med. Chem. Anal.* **2013**, *3*, 10; (b) Asif, M. *Int. J. Med. Chem.* **2014**, *2014*, 1; (c) Baltork, I. M.; Khosropour, A. R.; Moghadam, M.; Tangestaninejad, S.; Mirkhani, V.; Baghersad, S.; Mirjafari, A. C. *R. Chim.* **2011**, *14*, 944; (d) Selvam, T. P.; Kumar, P. V. *Res. Pharm.* **2011**, *1*, 1.
- [11] Bayouki, K. A. M. E.; Aly, M. M.; Mohamed, Y. A.; Basyouni, W. M.; Abbas, S. Y. *World J. Chem.* **2009**, *4*, 1619; (b) Obaid, A. M. A.; Hamide, S. G. A.; Kashef, H. A. E.; Aziz, A. A. M. A.; Azab, A. S. E.; Khamees, H. A. A.; Subbagh, H. I. E. *Eur. J. Med. Chem.* **2009**, *44*, 2379.
- [12] (a) Ma, J.; Li, P.; Li, X.; Shi, Q.; Wan, Z.; Hu, D.; Jin, L.; Song, B. *J. Agric. Food. Chem.* **2014**, *62*, 8928; (b) Wang, Z.; Wang, M.; Yao, X.; Li, Y.; Tan, J.; Wang, L.; Qiao, W.; Geng, Y.; Liu, Y.; Wang, Q. *Eur. J. Med. Chem.* **2012**, *53*, 275.
- [13] Noolvi, M. N.; Patel, H. M.; Bhardwaj, V.; Chauhan, A. *Eur. J. Med. Chem.* **2011**, *46*, 2327; (b) Zhang, G. H.; Xue, W. B.; An, Y. F.; Yuan, J. M.; Qin, J. K.; Pan, C. X.; Su, G. F. *Eur. J. Med. Chem.* **2015**, *95*, 377; (c) Kamal, A.; Bharathi, E. V.; Ramaiah, M. J.; Dastagiri, D.; Redd, J. S.; Viswanath, A.; Sultana, F.; Pushpavalli, S. N. C. V. L.; Bhadra, M. P.; Srivastava, H. K.; Sastry, G. N.; Juvekar, A.; Sen, S.; Zingde, S. *Bioorg. Med. Chem.* **2010**, *18*, 526.
- [14] Sharma, M.; Chauhan, K.; Shivahare, R.; Vishwakarma, P.; Suthar, M. K.; Sharma, A.; Gupta, S.; Saxena, J. K.; Lal, J.; Chandra, P.; Kumar, B.; Chauhan, P. M. S. *J. Med. Chem.* **2013**, *56*, 4374.
- [15] Pandey, S. K.; Singh, A.; Singh, A.; Nizamuddin. *Eur. J. Med. Chem.* **2009**, *44*, 1188; (b) Desai, N. C.; Vaghani, H. V.; Shihora, P. N. *J. Fluorine. Chem.* **2013**, *53*, 39.
- [16] Chandrika, P. M.; Rao, A. R. R.; Narsaiah, B.; Raju, M. B. *Int. J. Chem. Sci.* **2008**, *66*, 1119; (b) Nanthakumar, R.; Muthumani, P.; Girija, K. *Arabian. J. Chem.* **2014**, *53*, 1049.
- [17] (a) Lu, W.; Baig, I. A.; Sun, H. J.; Cui, C. J.; Guo, R.; Jung, I. P.; Wang, D.; Dong, M.; Yoon, M. Y.; Wang, J. G. *Eur. J. Med. Chem.* **2015**, *94*, 298; (b) Rashmi, A.; Ashish, K.; Gill, N. S.; Rana, A. C. *Int. J. Pharm.* **2011**, *2*, 22.
- [18] Havaladar, F. H.; Patil, A. R. *E-J. Chem.* **2008**, *5*, 347.

- [19] Wang, X.; Li, P.; Li, Z.; Yin, J.; He, M.; Xue, W.; Chen, Z.; Song, B. *J. Agric. Food. Chem.* **2013**, *61*, 9575.
- [20] Bouley, R.; Kumarasiri, M.; Peng, Z.; Otero, L. H.; Song, W.; Suckow, M. A.; Schroeder, V. A.; Wolter, W. R.; Lastochkin, E.; Antunes, N. T.; Pi, H.; Serge, V. A.; Vakulenko, S.; Hermoso, J. A.; Chang, M.; Mobashery, S. *J. Am. Chem. Soc.* **2015**, *137*, 17381.
- [21] Patil, D. A.; Patil, P. O.; Deshmukh, P. K.; Patil, G. B.; Shewale, B. D.; Patil, D. D.; Gattani, S. G. *Res. J. Pharm. Tech.* **2010**, *4*, 979.
- [22] Connolly, D. J.; Cusack, D.; Sullivan, T. P. O.; Guiry, P. J. *Tetrahedron* **2005**, *6*, 10153.
- [23] Mhaske, S. B.; Argade, N. P. *Tetrahedron* **2006**, *62*, 9787.
- [24] Mourad, A. E.; Aly, A. A.; Farag, H. H.; Beshr, E. A. *Beilstein. J. Org. Chem.* **2007**, *3*, 1.
- [25] Jiang, X.; Tang, T.; Chen, J. Z.; Zhu, Y.; Ji, S. *J. Org. Chem.* **2014**, *79*, 5082.
- [26] Zhou, J.; Fang, J. *J. Org. Chem.* **2011**, *76*, 7730.
- [27] Sharma, S.; Jain, A. *Tetrahedron Lett.* **2014**, *55*, 6051.
- [28] Wang, Y. F.; Zhang, F. L.; Chiba, S. *Org. Lett.* **2013**, *15*, 2842.
- [29] Yan, Y.; Xu, Y.; Niu, B.; Xie, H.; Liu, Y. *J. Org. Chem.* **2015**, *80*, 5581.
- [30] Wang, D.; Gao, F. *Chem. Cent. J.* **2013**, *7*, 1.
- [31] Heravi, M. M.; Derikvand, F.; Ranjbar, L. *Synth. Commun.* **2010**, *40*, 677.
- [32] Liu, J.; Lei, M.; Hu, L. *Green Chem.* **2012**, *14*, 840.
- [33] Chen, Z.; Chen, J. M.; Liu, J.; Ding, W.; Gao, X.; Wu, H. *J. Org. Chem.* **2013**, *78*, 11342.
- [34] Sahu, P. K.; Sahu, P. K.; Gupta, S. K.; Agarwal, D. D. *Ind. Eng. Chem. Res.* **2014**, *53*, 2085.
- [35] Kidwai, M.; Chauhan, R. *J. Mol. Catal. A: Chem.* **2013**, *377*, 1.
- [36] Sahu, P. K.; Sahu, P. K.; Jain, R.; Yadava, R.; Agarwal, D. D. *Catal. Sci. Technol.* **2012**, *2*, 2465.
- [37] Krishnamurthy, G.; Jagannath, K. V. *J. Chem. Sci.* **2013**, *125*, 807.
- [38] Heravi, M. M.; Ranjbar, L.; Derikvand, F.; Alimadadi, B.; Oskooie, H. H.; Bamoharram, F. F. *Mol. Divers.* **2008**, *12*, 181.
- [39] (a) Puttaraju, K. B.; Shivashankar, K.; Chandra, Mahendra, M.; Rasal, V. P.; Vivek, P. N. V.; Rai, K.; Chanu, M. B. *Eur. J. Med. Chem.* **2013**, *69*, 316; (b) Puttaraju, K. B.; Shivashankar, K. *RSC Adv.* **2013**, *3*, 20883; (c) Jagadishbabu, N.; Shivashankar, K. *RSC Adv.* **2015**, *5*, 95240.
- [40] Beerappa, M.; Shivashankar, K. *RSC Adv.* **2015**, *5*, 30364.
- [41] Godfrey, A. G.; Ganem, B. *Tetrahedron Lett.* **1990**, *31*, 4825.
- [42] Zheng, P.; Yan, L.; Ji, X.; Duan, X. *Synth. Commun.* **2010**, *41*, 16.
- [43] Adib, M.; Sheikhi, E.; Rezaei, N. *Tetrahedron Lett.* **2011**, *52*, 3191.
- [44] Keyume, A.; Esmayil, Z.; Wang, L.; Jun, F. *Tetrahedron* **2014**, *70*, 3976.
- [45] Khurana, J. M.; Nand, B.; Saluja, P. *Tetrahedron* **2010**, *66*, 5637.
- [46] Khumraksa, K.; Phakhodee, W.; Pattarawarapan, M. *Tetrahedron Lett.* **2013**, *54*, 1983.
- [47] Barbry, D.; Champagne, P. *Tetrahedron Lett.* **1996**, *37*, 7725.
- [48] Puligoundla, R. G.; Karnakanti, S.; Bantu, R.; Kommu, N.; Kondra, S. B.; Nagarapu, L. *Tetrahedron Lett.* **2013**, *54*, 2480.
- [49] Alinezhad, H.; Tajbakhsh, M.; Norouzi, M.; Baghery, S.; Akbari M. C. *R. Chim.* **2014**, *17*, 7.
- [50] Shao, Q.; Tu, S.; Li, C.; Cao, L.; Zhou, D.; Wang, Q.; Jiang, B.; Zhang, Y.; Hao, W. *J. Heterocycl. Chem.* **2008**, *45*, 411.
- [51] Yao, C.; Lei, S.; Wang, C.; Li, T.; Yu, C.; Wang, X.; Tu, S. *J. Heterocycl. Chem.* **2010**, *47*, 26.