Journal of Discrete Mathematical Sciences and Cryptography

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tdmc20>

Lict edge semientire graph of a planar graph

Y. B. Maralabhavi^a & Venkanagouda M. Goudar^a

^a Department of Mathematics , Bangalore University, Central College Campus ,
Bangalore , 560 001 , India

Published online: 03 Jun 2013.

To cite this article: Y. B. Maralabhavi & Venkanagouda M. Goudar (2007) Lict edge semientire graph of a planar graph, Journal of Discrete Mathematical Sciences and Cryptography, 10:3, 433-438, DOI: [10.1080/09720529.2007.10698130](https://doi.org/10.1080/09720529.2007.10698130)

To link to this article: <http://dx.doi.org/10.1080/09720529.2007.10698130>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Lict edge semientire graph of a planar graph

Y. B. Maralabhavi

Venkanagouda M. Goudar *

Department of Mathematics

Central College Campus

Bangalore University

Bangalore 560 001

India

Abstract

In this paper, we introduce the concept of the Lict edge semientire graph of a planar graph. We present characterizations of graphs whose lict edge semientire graphs are planar, outerplanar and Maximal outerplanar, crossing number one. Further, we establish a characterization of graphs whose lict edge semi entire graphs are Eularian and Hamiltonian.

Keywords and phrases : *Pathos, pathoslength, edge semientire graph, outerplanar, crossing number.*

1. Introduction

The lict graph⁽⁴⁾ $n(G)$ of a graph G is the graph whose vertex set is the union of the set of edges and the set of cutvertices of G in which two vertices are adjacent if and only if the corresponding edges of G are adjacent or the corresponding members of G are incident.

The edge semientire graph⁽³⁾ $e_e(G)$ of a plane graph G is the graph whose vertices can be put in one to one correspondence with the edges and regions of G in such a way that two vertices of $e_e(G)$ are adjacent if and only if the corresponding elements of G are adjacent. The line graph $L(G)$ of a graph G is the graph whose vertex set coincides with the edge set of G and in which two vertices are adjacent if and only if the corresponding edges are adjacent in G .

*E-mails: vmgouda@yahoo.co.in, v.goudar@rediffmail.com

Journal of Discrete Mathematical Sciences & Cryptography

Vol. 10 (2007), No. 3, pp. 433–438

© Taru Publications

A regionvertex is a vertex in lict edge semientire graph corresponding to the regions of G .

We now define the lict edge semientire graph $n_e(G)$ of a planar graph G whose vertex set is the union of the set of edges, set of vertices and set of regions of G in which two vertices are adjacent if and only if the corresponding edges are adjacent, edges are incident to the cutvertex and edges are lies on the region. In Figure 1, a graph G and its lict edge semientire graph $n_e(G)$ of a planar graph are shown.

Figure 1

All the undefined terms may be referred to Harrary². All graphs considered here are finite, undirected and without loops or multiple edges.

We need the following theorem for the proof of our further results.

Theorem 1. If G is a (p, q) graph whose vertices have degree d_i then $L(G)$ has q vertices and q_L edges where $q_L = -q + \frac{1}{2} \sum d_i^2$.

Theorem 2. The line graph $L(G)$ of a graph G has crossing number one if and only if G is planar and 1 or 2 holds:

- (1) The maximum degree $\Delta(G)$ is 4 and there is unique non cutvertex of degree.
- (2) The maximum degree $\Delta(G)$ is 5, every vertex of degree 4 is a cutvertex, there is a unique vertex of degree 5 and has almost 3 edges in any block.

Theorem 3. A connected graph G is isomorphic to its line graph if and only if it is a cycle.

Theorem 4. The lict graph $n(G)$ is planar if and only if $\deg v \leq 3$.

Theorem 5. The edge semi entire graph $e_e(G)$ is planar if and only if

- (1) G is a tree
- (2) $\deg v \leq 3$ for every vertex v of G .

Theorem 6. Let G be a plane graph. A necessary and sufficient condition for $e_e(G)$ to be eulerian is that each of the following holds:

- (1) Each edge of G is adjacent to even number of elements
- (2) Each region of G has even number of elements adjacent to it.

Theorem 7. If G is a hamiltonian plane graph, then $e_e(G)$ is also hamiltonian.

Lict edge semientire graph $n_e(G)$ of a planar graph G

We start with a few preliminary results.

Remark 1. For any graph G , $L(G) \subset n(G) \subset n_e(G)$ and $e_e(G) \subset n_e(G)$.

Remark 2. For any graph G , $n_e(G)$ is nonseparable.

Remark 3. If a graph G is K_2 then, $n_e(G)$ is complete.

Remark 4. If G itself is a block then $n_e(G) = e_e(G)$.

In the following theorem, we obtain the number of vertices and edges in lict edge semientire graph.

Theorem 8. For any (p, q) graph G whose vertices have degree d_i , cutvertices c , regions r and l_i be the number of edges to which cutvertex c_i belongs and e_k be the number of edges in which the region r_k lies, the lict edge semientire graph $n_e(G)$ has $(q + c + r)$ vertices and $\sum \left[\frac{d_i^2}{2} + l_i \right] + e_k$ edges.

Proof. By definition of $n_e(G)$, the number of vertices is the union of edges, cutvertices and regions of G . Hence $n_e(G)$ has $(q + c + r)$ vertices. Further the number of edges in $n(G)$ is the sum of number of edges incident with cutvertices in G , the sum of number of edges in $L(G)$. Since the number of edges in $L(G)$ has $-q + \sum \frac{d_i^2}{2}$, hence the number of edges in $n_e(G)$ is the sum of edges in $n(G)$ and the number of edges bounded by the regions e_k . In addition, the total number of edges that lie on the region $\sum r_i$ of G is q . Hence

$$\begin{aligned} E[n_e(G)] &= -q + \sum \left[\frac{d_i^2}{2} + l_i \right] + q + e_k \\ &= \sum \left[\frac{d_i^2}{2} + l_i \right] + e_k. \end{aligned}$$

Planar lict edge semientire graph

In this section, we obtain the condition for planarity of lict edge semientire graph.

Theorem 9. *The lict edge semientire graph $n_e(G)$ is planar if and only if $\deg v \leq 3$, for every vertex v of G .*

Proof. Suppose $n_e(G)$ is planar. Assume $\deg v \geq 4$. If there exists a vertex v of degree 4, then by definition, $L(G)$ is planar which contains $\langle K_4 \rangle$ as an induced subgraph. So the lict graph $n(G)$ contains $\langle K_5 \rangle$ as an induced subgraph, which is nonplanar. Clearly, $n_e(G)$ is also nonplanar, a contradiction.

Conversely suppose $\deg v \leq 3$. By the Theorem 5, $e_e(G)$ is planar and by the Theorem 4, $n(G)$ is also planar. Clearly, $n_e(G)$ is planar. $\square$

We now present a characterization of graphs whose lict edge semientire graph is outerplanar and maximum outerplanar.

Theorem 10. *The lict edge semientire graph $n_e(G)$ is outer planar if and only if G is a path.*

Proof. Suppose $n_e(G)$ is outerplanar. Assume that G has a vertex v of degree 3. The edges incident to v and the cutvertex v form $\langle K_4 \rangle$ as a subgraph in $n_e(G)$. Hence $n_e(G)$ is non outerplanar, a contradiction.

Conversely, suppose T is a path P_t of length $t \geq 1$. For $t = 1$, the result is obvious. For $t > 1$, the graph $n(G)$ has $(t - 1)$ blocks which are K_3 . Since G has exactly one region it follows that the region vertex, corresponding to this region, together with the above blocks form $(t - 1)$ number of induced subgraphs which are all $\langle K_4 - x \rangle$ in $n_e(G)$. Hence $n_e(G)$ is outerplanar. $\square$

Theorem 11. *The lict edge semientire graph $n_e(G)$ is maximum outerplanar if and only if G is a path.*

Proof. Proof follows from the Theorem 10. $\square$

In the next theorem, we characterize lict edge semientire graph in terms of crossing number one.

Theorem 12. *The lict semientire graph $n_e(G)$ has a crossing number one if and only if the following conditions hold:*

- (1) $\deg v \leq 3$, for every vertex v of G , and
- (2) G has unique vertex of degree 4, which is not a cutvertex.

Proof. Suppose $n_e(G)$ has crossing number one. Then it is nonplanar. By Theorem 9, $\deg v \geq 4$ for every vertex v of G . We now consider the following cases:

- Case 1.* Assume G has a vertex u of degree 5. If u is not a cutvertex, then by Theorem 4, the regionvertex is adjacent to edges of G . Clearly $C[n_e(G)] > 1$, a contradiction. If u is a cutvertex then the edges incident to this vertex together with the cutvertex form $\langle K_6 \rangle$ as a subgraph in $n(G)$, the regionvertex is adjacent to atleast one vertex of $\langle K_6 \rangle$ in $n_e(G)$. This gives $C[n_e(G)] > 1$, a contradiction.
- Case 2.* Assume G has atleast two vertices of degree 4. Suppose v_1 and v_2 are two noncutvertices of degree 4. Then $L(G)$ has atleast two crossings, by Theorem 2, $C[n_e(G)] > 1$, a contradiction. Suppose v_1 and v_2 are two cutvertices of degree 4. Then cutvertices v_1 and v_2 together with their corresponding four incident edges form two $\langle K_6 \rangle$ as subgraphs in $n(G)$ and hence in $n_e(G)$. Hence $C[n_e(G)] > 1$, a contradiction.

Conversely, suppose G holds both the conditions of the Theorem. Let v_1 be the noncutvertex of degree 4. Then by Theorems 2 and 3, $n(G)$ has crossing number one and hence $n_e(G)$ has crossing number one. $\square$

Theorem 13. *The lict edge semientire graph $n_e(G)$ is eulerian if and only if G is a cycle C_n , n is even.*

Proof. The cycle C_n does not contains a cutvertex. By the Remark 4, $n_e(G) = e_e(G)$. Also by Theorem 6, $e_e(G)$ is eulerian and hence $n_e(G)$ is eulerian.

Theorem 14. *The lict edge semientire graph $n_e(G)$ is hamiltonian if and only if $G \neq K_2$.*

Proof. Suppose $n_e(G)$ is hamiltonian. Assume that G is K_2 . Then this edge is incident with regionvertex w to form K_2 , which is nonhamiltonian, a contradiction.

Conversely, suppose $G \neq K_2$, we now consider the following cases:

- Case 1.* If G is a path and has exactly one regionvertex. Let $V[n(G)] = (e_1, e_2, \dots, e_n) \cup (c_1, c_2, \dots, c_{n-2})$, where $(c_1, c_2, \dots, c_{n-2})$ are cutvertices of G . Each block is a triangle and each block consist as vertices $B_1 = (e_1, c_1, e_2), B_2 = (e_2, c_2, e_3), \dots, B_n = (e_{n-1}, c_{n-2}, e_n)$. Also in $n_e(G)$, the regionvertex w is adjacent to $(e_1, e_2, \dots, e_n)$. Hence

$V[n_e(G)] = (e_1, e_2, \dots, e_n) \cup (c_1, c_2, \dots, c_{n-2}) \cup w$ form a cycle $we_1c_1e_2c_2e_3 \dots e_{n-1}c_{n-1}c_nw$ containing all the vertices of $n_e(G)$. Clearly $n_e(G)$ is hamiltonian.

Case 2. If G is a tree and has exactly one regionvertex. Let $[n(G)] = (e_1, e_2, \dots, e_n) \cup (c_1, c_2, \dots, c_j)$, where $(c_1, c_2, \dots, c_j)$ are the cutvertices of G . Clearly, each block is K_3 if degree of the cutvertex is two and is K_4 if degree of the cutvertex is three. In $n_e(G)$, the regionvertex w is adjacent to $(c_1, c_2, \dots, c_j)$. By Remark 2, $n_e(G)$ is nonseparable. Clearly, the vertices $(e_1, e_2, \dots, e_n) \cup (c_1, c_2, \dots, c_j) \cup w$ form $we_1c_1e_2c_2e_3e_4 \dots c_jc_nw$ containing all the vertices of $n_e(G)$. Hence $n_e(G)$ is hamiltonian.

Case 3. If G is hamiltonian graph, then by Theorem 7, $e_e(G)$ is hamiltonian. Hence $n_e(G)$ is hamiltonian.

Case 4. If G is the graph other than above types of graphs, then by Remark 2, $n_e(G)$ is nonseparable, hence it is hamiltonian.

References

- [1] F. Harrary (1977), Covering and packing in graphs – I, *Annals of New York Academy of Science*, 175:198.
- [2] F. Harrary (1969), *Graph Theory*, Addison-Wesley, Reading Mass., p. 72, 107.
- [3] V. R. Kulli and D. G. Akka (1980), On semientire graphs, *Journal Mathematical Science*, Vol. 14 (6), pp. 585–588.
- [4] M. H. Muddebihal (1991), *Some Results in the Theory of Graphs*, Ph.D. Thesis, Gulbarga University, Gulbarga, India.
- [5] J. Sedlacek (1963), Some properties of interchange graphs, in *Proceeding of the Symposium Smolenics*, Academic Press, New York, p. 145.

Received August, 2006